

## Iron

NO. 12

### NORTH AMERICAN EDITION

**I**ron (Fe) is a nutrient required by all organisms, including microbes, plants, animals, and humans. It was first recognized as a necessary plant nutrient in the mid 19th century when Fe-deficient grapes were successfully treated with foliar applications of Fe salts. Iron is a component of many vital plant enzymes and is required for a wide range of biological functions. It is common in the earth's crust and as a result, most soils contain abundant Fe, but in forms that are low in solubility and sometimes not readily available for plant uptake.

#### Iron in Soils

Iron is abundant in many rocks and minerals and as soils develop there can be either enrichment or depletion of Fe. Depletion commonly leads to deficiency and enrichment can cause toxicity in unique conditions. The main source of Fe in soils for use by plants comes from secondary oxide minerals that are adsorbed or precipitated onto soil mineral particles and organic matter. Although Fe is very abundant, its availability for plant uptake is quite low.

#### Iron in Plants

Plant roots absorb Fe from the soil solution most readily as (ferrous)  $Fe^{2+}$  but in some cases also as (ferric)  $Fe^{3+}$  ions. The chemical nature of Fe allows it to play an essential role in oxidation and reduction reactions, respiration, photosynthesis, and enzyme reactions. For example, Fe is an important component of the enzymes used by nitrogen-fixing bacteria.

The Fe concentration in plant leaf tissues varies between plant species, but is generally between 50 and 250 ppm (dry weight basis). If the Fe concentration is less than 50 ppm there are usually signs of deficiency, and toxic effects may be observed when the concentration exceeds 500 ppm.

The solubility of Fe oxide minerals in soil is very low, so plant roots have two general ways to access the  $Fe^{2+}$  or  $Fe^{3+}$  ions. The first strategy occurs in dicot species, and non-grass monocot species where  $Fe^{3+}$  ions are reduced to  $Fe^{2+}$  ions before moving into the root across selective membranes. This process involves the root excreting a variety of organic compounds and acids into the soil. In the second strategy, roots of grass species acquire Fe by excreting an organic chelate (siderophore) that solubilizes Fe from the soil, allowing enhanced uptake.



IPNI IMAGE/BIANCHINI



IPNI IMAGE/SHARWA AND KUMAR



IPNI IMAGE/SHARWA AND KUMAR

Iron deficiency in soybean (left), sorghum (middle), and wheat (right).

#### Soil Factors and Iron Deficiency

Most soils contain adequate Fe for plant nutrition, but chemical and environmental factors restrict plant uptake. Iron deficiencies are commonly observed in soils with elevated pH (>7.5), especially where there is abundant calcium carbonate (lime). Iron solubility is greatly increased as soil pH drops into the acidic range.

Soils containing abundant calcium carbonate can form bicarbonate ions ( $HCO_3^-$ ) if the soils become overly wet, and bicarbonate interferes with Fe uptake by plants. This inhibition is usually only temporary and Fe deficiency symptoms disappear when the soil drains and warms up.

When soils become saturated,  $Fe^{3+}$  becomes converted to  $Fe^{2+}$  by microbial action. The  $Fe^{2+}$  form is much more soluble and can even result in toxicity for some rice varieties in flooded soils under strongly acid conditions.



INTERNATIONAL  
PLANT NUTRITION  
INSTITUTE

3500 PARKWAY LANE, SUITE 550  
PEACHTREE CORNERS, GEORGIA 30092-2844 USA  
PHONE (770) 447-0335 | WWW.IPNI.NET

Plants growing in soils with low organic matter content are generally more susceptible to Fe deficiency than with abundant organic matter. Humus compounds are effective at binding and releasing Fe ions into soil solution. Portions of a field that are eroded (low soil organic matter) tend to be more susceptible to Fe deficiency.

Since many soil and environmental factors combine to regulate the Fe supply to plants, there is no widely accepted method of testing soils to predict the need for supplemental fertilization.

### Deficiency and Toxicity Symptoms

Iron deficiency symptoms are universal among plant species, with general stunting and yellowing of younger leaves. Young Fe-deficient leaves develop chlorosis (yellowing) between the leaf veins, while the veins initially remain green. As the deficiency becomes more severe, the younger leaves become pale yellow to white in color. The young tissue is impacted first because Fe is poorly mobile within plants and does not readily translocate from older to younger tissues.

Iron toxicity is relatively rare, but the symptoms include bronzed and striped leaves. These effects are the result of

excess Fe-hydroxyl radicals disrupting cellular functions. Due to the importance of maintaining Fe concentrations within safe ranges in plant tissues, the whole process of Fe uptake into roots (i.e., the movement from roots to shoots, and storage and release within plant cells) is highly regulated.

Tissue analysis for Fe can be complicated since any dust that may be present on the leaf surface will also contain Fe. Rinsing or washing plant leaves is recommended prior to Fe analysis. Most tissue analyses rely on sampling the young leaves, since they are generally the first to show deficiency symptoms.


### Fertilizing for Iron Deficiency

When inorganic Fe fertilizers are added to soil (e.g., ferric sulfate, ferrous sulfate, ferrous ammonium phosphate, ferrous ammonium sulfate, and oxides of Fe), they are rapidly converted to insoluble forms and provide minimal benefit for plant nutrition. Iron fertilizers protected with an organic chelate can be effectively applied to soils to correct plant deficiencies. For example, chelated fertilizers such as Fe-EDDHA and Fe-EDTA have been used with reasonable effectiveness (Table 1), but their cost is often prohibitive for large-scale application. Foliar sprays containing Fe salts or chelates are effective at correcting plant Fe deficiencies during the growing season, but they may require repeated applications to prevent reoccurrence of deficiency.

### Crop Response

Several remedies are used to compensate for plant Fe deficiency. Depending on local conditions, some of these solutions may be more practical than others.

- Grow plant varieties and cultivars specifically adapted to the local conditions that are tolerant of low-Fe conditions. Large genetic differences exist among cultivars and a variety change is often effective for dealing with challenging soil conditions. (Figure 1).



**Figure 1.** Grain yield of three soybean varieties grown in calcareous soil. The Glacier variety, susceptible to Fe deficiency, responded to foliar Fe application while no yield response was observed for more tolerant Council and Traill varieties<sup>1</sup>. Error bars denote an LSD (0.05).

- Apply a Fe-containing fertilizer in the form of an inorganic salt or a chelated material to the soil.
- Spray a Fe-containing solution onto plant leaves to prevent or correct deficiencies. This does not correct any underlying soil problems preventing uptake of adequate Fe, but it can assist with eliminating growth limitations from Fe deficiency.
- Add an acidifying material to soils with elevated pH to improve the solubility of Fe. This acidification can be done for the entire field or spot treatment of a portion of the root zone is often sufficient to improve Fe availability.

- Improve Fe availability by growing two plant species together. The ability of one crop to solubilize and acquire Fe sometimes results in sharing with a companion crop that has lesser capacity to extract Fe (Table 1).

**Table 1.** Relative grain yield of two soybean varieties, one susceptible and one tolerant to Fe deficiency, compared with the tolerant variety grown on a Fe-sufficient site (100%)<sup>2</sup>.

Oat companion crop	Fe-chelate fertilizer	Fe chlorosis susceptible variety	Fe chlorosis tolerant variety
Relative yield, %			
No	No	48 e	82 c
No	Yes	71 d	87 bc
Yes	No	73 d	76 cd
Yes	Yes	87 bc	93 ab

Treatments include addition of a Fe-chelate fertilizer, or the presence of an oat companion crop on severely Fe-deficient soils. Letters following relative yields indicate significance at  $p \leq 0.10$  for both varieties.

### References

1. Goos, R.J. and B.E. Johnson. 2000. *Agron. J.* 92:1135-1139.
2. Kaiser, D.E., J.A. Lamb, P.R. Bloom, and J.A. Hernandez. 2014. *Agron. J.* 106:1963-1974.